

AD627368

AD

US ARMY MEDICAL RESEARCH LABORATORY

FORT KNOX, KENTUCKY

REPORT NO. 640

THE EFFECT OF pH, TEMPERATURE, ANTIVENIN AND
FUNCTIONAL GROUP INHIBITORS ON THE TOXICITY AND
ENZYMATIC ACTIVITIES OF CROTALUS ATROX VENOM

by

John H. Brown, Ph. D.

20 September 1965

Code 1

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION		
Hardcopy	Microfiche	
\$ 100	\$ 050	22 PP
ARCHIVE COPY		



UNITED STATES ARMY
MEDICAL RESEARCH AND DEVELOPMENT COMMAND

20050307000

Defense Documentation Center Availability Notice

Qualified requesters may obtain copies of this report from
DDC.

Disposition

Destroy this report when it is no longer needed. Do not return
it to the originator.

Disclaimer

The findings in this report are not to be construed as an official
Department of the Army position, unless so designated by other author-
ized documents.

Acknowledgement

The author wishes to express his appreciation to M. Edith
Bowles, Pfc Norman L. Schwartz and Pfc Donald J. Cherland for
their technical assistance.

In conducting the research described in this report,
the investigators adhered to the "Principles of Labora-
tory Animal Care as established by the National Society
for Medical Research."

83E:28C:4

ABSTRACT

THE EFFECT OF pH, TEMPERATURE, ANTIVENIN AND FUNCTIONAL GROUP INHIBITORS ON THE TOXICITY AND ENZYMATIC ACTIVITIES OF CROTALUS ATROX VENOM

OBJECT

To determine the effects of certain group-specific reagents on the toxicity of Crotalus atrox (Western diamondback rattlesnake) venom and the effect of pH, temperature, and antivenin on the toxicity and enzymatic activities of this venom.

RESULTS

Crotalus atrox venom has been completely detoxified by 50 mM N-bromosuccinimide. This venom cannot be detoxified with the sulfhydryl group reagents, N-ethylmaleimide or iodoacetamide, or the metal-binding agent EDTA. It is very slightly detoxified by reaction with O-methylisourea, diisopropylfluorophosphate, or British anti-lewisite. The toxic principle in C. atrox venom is resistant to low pH treatment, but is destroyed by heating at 75° for one hour. Under the same conditions the following enzymes are destroyed: phosphodiesterase, esterase (TAMEase), and proteolytic enzyme activity on azocoll, congocoll, casein, and N-benzoyl-DL-arginine p-nitroanilide. Phospholipase A activity is only partially destroyed under these conditions.

CONCLUSIONS

The effects observed on various enzymes with functional group inhibitors and the results of the heat inactivation and acid treatment of C. atrox venom indicate that the main toxicity of this venom is not associated with the enzymatic components but may reside with a peptide of relatively low molecular weight.

RECOMMENDATIONS

Attempts should be made to separate the proteolytic, phospholipase A, anticoagulant and toxic activities of C. atrox venom and

their physiological effects determined in order to learn whether the toxicity of this venom is due to separated toxin(s) or a synergistic action of one or more enzymes. Detoxification with N-bromosuccinimide and antibody production should be studied with the detoxified venom produced.

TABLE OF CONTENTS

	<u>Page No.</u>
INTRODUCTION	1
MATERIALS AND METHODS	2
RESULTS	2
DISCUSSION	6
SUMMARY	9
REFERENCES	10
 Table 1	 3
Table 2	4
Tables 3 and 4	5
Table 5	6
Table 6 and Figure 1	7

THE EFFECT OF pH, TEMPERATURE, ANTIVENIN AND FUNCTIONAL GROUP INHIBITORS ON THE TOXICITY AND ENZYMATIC ACTIVITIES OF CROTALUS ATROX VENOM

INTRODUCTION

Various chemical compounds have been used to detoxify snake venoms. Trethewie (1) found that tetra-hydro-aminoacridine and poly-vinyl-pyrrolidone increases the survival time of mice following injection of cobra venom in vitro. Heparin and other sulphated mucopolysaccharides can inhibit the lethal effects of Russell's viper venom (Vipera russellii) (2). Also unsaturated fatty acids can protect mice against Crotalus terrificus venom (3). Hydro-cortisone, in large doses will ameliorate the lethality of cobra venom in dogs (4). Deichmann et al. (5) found similar results with hydro-cortisone on Crotalus adamanteus venom. Sawai et al. (6) have shown that treatment of snakebite with ethylenediaminetetraacetic acid (EDTA) enhances the effect of antivenin in Habu (Trimeresurus flavoviridis) bite. Both the necrotic and lethal effects of Habu venom have been partially neutralized by dihydrolipoic acid and tetracycline in vivo (7). The mechanisms by which these compounds alter the toxicity of snake venoms is unknown.

Kochwa et al. (8) showed that horse or rabbit antivenin prepared from whole Vipera xanthina palestinae venom will not neutralize the neurotoxic components of that venom but will neutralize the hemorrhagic activity. Pooled sera of rabbits immunized with either Walterinnesia aegyptia or Pseudocerastes fieldii venom inactivated the venom phospholipase activities (9). Rabbit antivenin for Echis colorata venom was capable of neutralizing proteolytic activity, but not the amino acid oxidase, phospholipase, erythrocyte spherizing factor, the anticoagulant, the procoagulant or the fibrinogenolysin-fibrinolysin activity (10). H β -Proteinase of Trimeresurus flavoviridis (Habu) venom is completely inhibited by anti-Habu serum (11). It is possible that some of the antigens which combine with snake venom antivenin play no significant role in the toxicity of the venom.

This study describes the effects of certain group-specific reagents on the toxicity of Crotalus atrox venom, the effect of pH, temperature, and Antivenin Crotalidae Polyvalent (Wyeth Laboratories) on the toxicity and enzymatic activities of this venom. Evidence is also presented which indicates that one enzyme of C. atrox venom has both esterolytic and proteolytic activities.

MATERIALS AND METHODS

Venom of C. atrox was collected from several hundred rattlesnakes, immediately frozen, and lyophilized within two weeks after collection. It was sieved and stored in the cold in a desiccator. Immediately before use the venom was reconstituted by dissolving in distilled water or appropriate solvent and centrifuged to remove undissolved matter, if any.

Studies on toxicity of lyophilized C. atrox venom were carried out in 20 g albino mice by intraperitoneal injection of 0.5 ml of venom solutions. Four mice were used for each of ten doses: 50, 62, 73, 80, 89, 100, 114, 160, 200, and 400 μ g per mouse. The LD₅₀ was estimated from a log dose-probit curve to be 112 γ of lyophilized venom per mouse.

Enzyme activities were measured as previously described (12-14). Venom and antivenin were incubated at 37°C for 60 minutes in 0.025 M Tris-HCl (tris-(hydroxymethyl)aminomethane), pH 8.0 in various proportions. Aliquots were withdrawn, diluted appropriately, and the enzymic activities measured. Controls without antivenin and without venom were determined with the venom-antivenin mixtures at the same time to standardize venom-enzyme activities and to determine whether antivenin had any enzyme activity.

All chemicals used were reagent grade.

RESULTS

Table 1 lists the effects of several functional group reagents on the toxicity of C. atrox venom. O-Methylisourea (OMIU), diisopropylfluorophosphate (DFP), and British anti-lewisite (BAL) (2,3-dimercaptopropanol) showed a slight inhibition of toxicity. With BAL, however, a precipitate was formed which could indicate a denaturation of the toxin(s). Only with 50 mM N-bromosuccinimide (NBS) was there complete detoxification of the venom. This compound, NBS, has been shown to react fairly specifically with tryptophan residues of proteins (15,16). N-Ethylmaleimide (NEM), EDTA, and iodoacetamide (IAA) had little effect on toxicity.

The toxic principle in C. atrox venom is quite resistant to treatment at low pH values, as shown in Table 2. Venom treated with 0.10 N HCl for one hour at 37°C was still toxic.

TABLE 1

EFFECT OF FUNCTIONAL GROUP INHIBITORS ON THE TOXICITY
OF CROTALUS ATROX VENOM

Venom (11.25 mg/ml) was incubated at 37°C for one hour with the concentrations of inhibitors listed then dialyzed overnight in the cold with one or two changes of distilled water. One-half ml of each of these solutions, after dilution containing the number of LD₅₀ listed was injected into 20 g albino mice intraperitoneally. With the exception of DFP and NBS, which were incubated in 0.1 M sodium phosphate, pH 8.0, the compounds were incubated with venom in 0.05 M Tris-HCl, pH 8.0. The fractions in parentheses indicate the number of deaths over the number of mice injected.

Inhibitor	Conc. mM	LD ₅₀ Injected No.	Survival %
OMIU	*	5	20 (4/5)
"	*	10	0 (5/5)
EDTA	1	5	0 (4/4)
NEM	100	5	0 (5/5)
IAA	100	5	0 (5/5)
DFP	50	5	20 (4/5)
"	50	10	0 (5/5)
NBS	10	10	0 (5/5)
"	50	150	100 (0/5)
BAL	20	5**	100 (0/5)
"	20	10**	0 (5/5)

*Venom (11.25 mg/ml) was incubated for 72 hours at 2°C with a 50:1 excess of O-methylisourea (OMIU). The unreacted OMIU was removed by dialysis at 2°C against distilled water. The OMIU-treated venom was then incubated at 37°C and samples were removed for assay after 60 minutes.

**Slight precipitate noted.

TABLE 2

EFFECT OF pH ON THE TOXICITY OF CROTALUS ATROX VENOM

Venom (4 LD₅₀/ml) was treated for 15, 30, and 60 minutes at 37°C in the following solvents: pH 1.0, 0.10 N HCl; pH 3.0, 0.05 M glycine; pH 4.0, 5.0, 6.0, 0.05 M sodium acetate; pH 8.0, 0.05 M Tris-HCl. Following incubation, 0.5 ml of venom (in the respective solvent) was injected. Controls contained no venom. Numbers in parentheses are deaths over the total number of mice injected.

LD ₅₀ Injected No.	Incubation Time Min.	pH					
		1	3	4	5	6	8*
2	15	(3/3)	(3/3)	(2/3)	(3/3)	(3/3)	-
2	30	(3/3)	(2/3)	(3/3)	(3/3)	(3/3)	-
2	60	(3/3)	(3/3)	(3/3)	(2/3)	(3/3)	(3/3)
0							
0	60	(0/3)**	(0/3)	(0/3)	(0/3)	(0/3)	(0/3)

*Treated with 0.05 M Tris-HCl, pH 8.0, at 25°C.

**These controls died about four days after injection.

The toxic principle in C. atrox venom is not very resistant to heat, as shown in Table 3. Treating the venom at 75°C for one hour destroys most of the toxicity and treatment at 100°C for the same time completely inactivates the toxic component(s).

Table 4 lists the effect of heat on the enzymes in C. atrox venom. Most of the enzymes are inactivated at temperatures between 50° and 75°C with the exception of phospholipase A which retains more than half of its activity after heating it for one hour at 100°C. Also, 17% of the proteolytic activity on benzoyl-DL-arginine p-nitroanilide (BAPNA) is retained after treatment at 75°C for one hour.

Table 5 lists the effect of antivenin on the hydrolysis of azocoll, BAPNA, casein and congocoll by C. atrox venom. With a 10:1 ratio (w/w) of antivenin:venom only slight inhibition was observed. Higher ratios (50:1, 100:1), however, produced significant inhibition. There is an obvious difference in the effect of antivenin on the hydrolysis of

TABLE 3

**EFFECT OF TEMPERATURE ON THE TOXICITY OF
CROTALUS ATROX VENOM**

Venom (11.25 mg/ml) was treated in 0.05 M Tris-HCl, pH 8.0, at 0°, 25°, 50°, 75°, and 100°C for 60 minutes then, after dilution, 0.5 ml was injected into mice. Numerators are deaths; denominators are total of mice injected.

LD ₅₀ Injected No.	0°	25°	50°	75°	100°
2	3/3	3/3	3/3	0/3*	0/3*
5	-	-	-	0/3	0/3
10	-	-	-	0/3	0/3
15	-	-	-	1/8	0/3
20	-	-	-	5/8	0/5
30	-	-	-	7/8	0/5
50	-	-	-	7/8	0/8

*Precipitates were formed at these temperatures. The supernatant liquid obtained after centrifugation, contained 65% of the biuret positive material. This was used for injection.

TABLE 4

**EFFECT OF TEMPERATURE ON THE ENZYMATIC ACTIVITIES
OF CROTALUS ATROX VENOM**

Venom (11.25 mg/ml) was incubated in 0.05 M Tris-HCl, pH 8.0, at 0°, 25°, 50°, 75°, and 100°C for 60 minutes and then diluted appropriately for assay. Enzymic assays were carried out as described previously (12-14).

Temperature	0	25	50	75*	100*
	Per Cent Activity				
Proteolytic Substrates					
Asocoll	100	100	95	0	0
Congocoll	100	119	81	0	0
BAPNA	100	85	79	17	0
Casein	100	80	31	0	0
Esterases					
Phosphodiesterase	100	100	49	0	0
Esterase (TAMEase)	100	107	77	0	0
Phospholipase A	100	-	-	67	58

*Precipitates were formed at these temperatures. The supernate, obtained after centrifugation, contained 65% of the biuret reading material. This was used for assaying the enzymes.

TABLE 5

**EFFECT OF ANTIVENIN ON THE PROTEOLYTIC ENZYME
ACTIVITIES OF CROTALUS ATROX VENOM**

Substrate	mg Antivenin mg Venom	Per Cent Inhibition
Azocoll	50/5.0	12
"	50/0.5	92
BAPNA	50/5.0	9
"	50/1.0	27
Casein	50/5.0	26
"	50/1.0	57
Congocoll	50/5.0	16
"	50/0.5	100

BAPNA and casein by C. atrox venom. Hydrolysis of BAPNA is not inhibited appreciably, while caseinolytic activity is inhibited almost 60 per cent at the 50:1 ratio of antivenin:venom.

Table 6 lists the effect of antivenin on the esterase activities of C. atrox venom. At a low ratio of antivenin to venom (10:1), phosphodiesterase is markedly inhibited while hydrolysis of TAME (N-p-toluenesulphonyl-L-arginine methyl ester) is only slightly inhibited by a 17:1 ratio of antivenin to venom. Phospholipase A activity was not inhibited by a 10:1 ratio of antivenin to venom.

Figure 1 shows the effect of diisopropylfluorophosphate (DFP) on the hydrolysis of TAME and BAPNA by C. atrox venom. The hydrolytic activity on TAME or BAPNA was essentially equally inhibited at any concentration of DFP used. It thus appears that C. atrox venom contains an enzyme which can hydrolyze both — CONH — and — COOR — bonds, viz. amide and ester linkages.

DISCUSSION

The toxic principle in C. atrox venom has been shown to be quite resistant to treatment at low pH values. If the toxic principle

TABLE 6

**EFFECT OF ANTIVENIN ON THE ESTERASE ACTIVITIES
OF CROTALUS ATROX VENOM**

For experimental details see Materials and Methods.

Esterase	mg Antivenin mg Venom	Per Cent Inhibition
Phosphodiesterase	50/5.0	86
Esterase	50/5.0	18
"	50/3.0	24
Phospholipase A	50/5.0	0




Fig. 1. Effect of diisopropylfluorophosphate on the hydrolysis of BAPNA and TAME by Crotalus atrox venom. Venom (6.0 mg/ml) and DFP were incubated for 60 minutes in 0.05 M Tris-HCl buffer, pH 8.0, at 37°C. Aliquots were removed, diluted with water to 1.0 mg/ml and assayed for enzymic activity.

were a low-molecular weight peptide, it could be non-denaturable at low pH values simply because it has little secondary and tertiary structure to be altered. Yang (17) has recently shown that more than four-fifths of Russell's viper and Formosan Habu venoms are composed of components with molecular weights less than 10,000. He also demonstrated that about one-third of Green Habu (Trimeresurus gramineus formensis), Hundred-pace snake (Agkistrodon acutus), and Eastern diamondback (C. adamanteus) venoms are composed from protein components larger than 50,000; more than 50% of them were low-molecular weight proteins ranging 4,500 - 10,000. Preliminary separation studies* on C. atrox venom with Bio-Gel (P-60) indicate that separated and lyophilized components with molecular weights 5,000 - 14,000 have about twice the toxicity as whole lyophilized venom in terms of either protein or dry weight. These low-molecular weight components constitute approximately 43% of the dry venom. This increase in toxicity is paralleled by increases in proteolytic enzyme activities and phosphodiesterase activity.

Previous work in this laboratory (12-14) demonstrated that most of the enzymes in C. atrox venom could be almost completely inactivated by one or more of the functional-group inhibitors listed in Table 1, which have little or no effect on the toxicity of this venom. One millimolar EDTA, although it has little effect on the toxicity, completely inhibits the proteolytic activity of C. atrox venom on casein, azocoll, and congocoll, as well as phosphodiesterase activity. Ten millimolar NBS has been shown to inhibit 94% of the proteolytic activity on BAPNA, yet this does not affect toxicity. Esterase (TAMEase) activity is completely inhibited by DFP; this compound has little effect on toxicity. From these observations and from the data in this paper, one can presume that the toxic principle in C. atrox venom is not associated with any of the enzymes listed in Table 4.

Of the functional group reagents used in an attempt to detoxify C. atrox venom only NBS was completely effective in reducing the toxicity to nil. This compound has been used by Boroff et al. (18) to detoxify Clostridium botulinum toxin. It has also been used by Okada et al. (19) to inactivate Bacillus subtilis α -amylase with no loss of antigenicity and loss of one mole per mole of tryptophan. In preliminary experiments with the NBS-detoxified C. atrox venom there was some indication that the treated venom could elicit antibody formation.

L-Amino acid oxidase and adenosine triphosphatase, which are present in C. atrox venom (20, 21), have not, however, been characterized with respect to these functional-group inhibitors. Other enzymes

*Unpublished results.

which have not been investigated and could be present in this venom, based on their presence in other crotalid venoms, include deoxyribonuclease (DNase), ribonuclease (RNase), diphosphopyridine nucleotide-pyrophosphatase, hyaluronidase and 5'-nucleotidase. We have observed that C. atrox venom has little, if any, hyaluronidase activity,* using the colorimetric method of Slotta and Ballester (22).

Furthermore, phospholipase A activity is only partially inactivated by heating venom at 100°C, whereas the toxicity is completely destroyed under these conditions.

The effect that antivenin has on the enzymes of C. atrox venom, under the conditions listed, indicates that phospholipase A has little antigenicity. The enzymes responsible for the hydrolysis of TAME, BAPNA, and casein are weakly antigenic and phosphodiesterase, azocollase, and congocollase are strongly antigenic.

Some of these enzymic activities in this and other venoms have been shown to be toxic per se (23-27). Whether the toxicity of this venom is due to separate toxin(s) or the synergistic action of several enzymes has yet to be determined.

Mandelbaum and Henriques (28) have shown the Bothrops protease A of Bothrops jararaca venom is the enzyme responsible for the hydrolysis of N-benzoyl-L-arginine amide (BAA) and that this crude enzyme also contains the bulk of TAMEase activity. It may be that one enzyme is hydrolyzing both TAME and BAA. Evidence in this paper (Fig. 1) indicates that one enzyme might have both proteolytic and esterolytic activity similar to chymotrypsin and trypsin (29).

SUMMARY

1. C. atrox venom has been completely detoxified by 50 mM N-bromosuccinimide.

2. This venom cannot be detoxified by reaction with the sulfhydryl group reagents, N-ethylmaleimide or iodoacetamide, or the metal-binding agent EDTA.

3. C. atrox venom is very slightly detoxified by reaction with O-methylisourea, diisopropylfluorophosphate, or British anti-lewisite.

*Unpublished results.

4. The toxic principle in C. atrox is resistant to low pH treatment, but is destroyed by heating at 75° or 100°C for one hour. Under the same conditions the following enzymes are destroyed: phosphodiesterase, esterase (TAMEase), and proteolytic enzyme activity on azocoll, congocoll, casein, and N-benzoyl-DL-arginine p-nitroanilide. Phospholipase A activity is only partially destroyed under these conditions.

5. Polyvalent crotalid antivenin has been shown to inhibit those enzymic activities of C. atrox venom responsible for the hydrolysis of TAME, BAPNA, casein, azocoll, and congocoll. Phosphodiesterase was significantly inhibited by antivenin but phospholipase A, under the conditions studied, was not inhibited.

6. Diisopropylfluorophosphate was found to inhibit the hydrolysis of TAME and BAPNA by C. atrox venom almost identically over a wide range of DFP concentrations.

REFERENCES

1. Trethewie, E. R. The effect of poly-vinyl-pyrrolidone and tetra-hydro-aminoacridine on the mortality and survival time of mice injected with snake venom. Med. J. Aust. 43: 8, 1956.
2. Higginbotham, R. D. Sulfated polysaccharides and resistance to Russell's viper venom. Proc. Soc. Exper. Biol. Med. 110: 135, 1962.
3. Spink, W. W. and C. K. Su. Comparative protective action of unsaturated fatty acids for mice against exotoxin, endotoxin and snake venom. Proc. Soc. Exper. Biol. Med. 112: 463, 1963.
4. Morales, F., H. D. Root, and J. F. Perry, Jr. Protection against Naja naja venom in dogs by hydrocortisone. Proc. Soc. Exper. Biol. Med. 108: 522, 1961.
5. Deichmann, W. B., J. L. Radomski, J. J. Farrell, W. E. MacDonald, and M. L. Keplinger. Acute toxicity and treatment of intoxications due to Crotalus adamanteus (rattlesnake venom). Amer. J. Med. Sci. 236: 204, 1958.

6. Sawai, Y., M. Makino, I. Tatenno, T. Okonogi, and S. Mitsuhashi. Studies on the improvement of treatment of Habu snake (Trimeresurus flavoviridis) bite. 3. Clinical analysis and medical treatment of Habu snake bite on the Amami Islands. Jap. J. Exptl. Med. 32: 117, 1962.
7. Sawai, Y., M. Makino, and Y. Kawamura. Studies on the antitoxic action of dihydrolipoic acid (dihydrothiopic acid) and tetracycline against Habu snake (Trimeresurus flavoviridis HALLOWELL) venom. In Venomous and Poisonous Animals and Noxious Plants of the Pacific Area, H. L. Keegan and W. V. MacFarlane (eds.), Pergamon Press, New York, 1963, pp. 327-335.
8. Kochwa, S., S. Gitter, A. Strauss, A. De Vries and M. Leffkowitz. Immunologic study of Vipera xanthina palestinae venom and preparation of potent antivenin in rabbits. J. Immunol. 82: 107, 1959.
9. Gitter, S., C. Moroz-Perlmutter, J. H. Boss, E. Livni, J. Rechnic, N. Goldblum and A. De Vries. Studies on the snake venoms of the Near East: Walterinnesia aegyptia and Pseudocerastes fieldii. Am. J. Trop. Med. Hyg. 11: 861, 1962.
10. Gitter, S., G. Levi, S. Kochwa, A. De Vries, J. Rechnic and J. Casper. Studies on the venom of Echis colorata. Am. J. Trop. Med. Hyg. 9: 391, 1960.
11. Maeno, H. and S. Mitsuhashi. Studies on H β -proteinase of Habu snake venom with special reference to substrate specificity and inhibitory effect of serum. J. Biochem. 50: 330, 1961.
12. Brown, J. H. Effect of enzymatic functional group inhibitors on proteolytic enzymes of Crotalus atrox venom. USAMRL Report No. 622, Fort Knox, Ky., 1965.
13. Brown, J. H. and M. Edith Bowles. Effect of enzymatic functional group inhibitors on the esterases of Crotalus atrox venom. USAMRL Report No. 627, Fort Knox, Ky., 1965.

14. Brown, J. H. and M. Edith Bowles. Studies on the phospholipase A activity of Crotalus atrox venom. USAMRL Report No. 630, Fort Knox, Ky., 1965.
15. Viswanatha, T. and W. B. Lawson. The action of N-bromosuccinimide on chymotrypsin. Arch. Biochem. Biophys. 93: 123, 1961.
16. Viswanatha, T., W. B. Lawson, and B. Witkop. The action of N-bromosuccinimide on trypsinogen and its derivatives. Biochim. Biophys. Acta, 40: 216, 1960.
17. Yang, C. Fractionation of snake venom on Sephadex. J. Formosan Med. Assn. 62: 611, 1963.
18. Boroff, D. A., B. R. Dasgupta, and R. J. Suhadolnik. Chemistry and biological activity of Clostridium botulinum. Biochem. Pharmacol. 12 (Abstract Volume): 228, 1963.
19. Okada, Y., K. Onoue, S. Nakashima, and Y. Yamamura. Studies on enzyme-antienzyme systems. II. N-Bromosuccinimide modified bacterial- α -amylase. J. Biochem. 54: 477, 1963.
20. Kearney, E. B. and T. P. Singer. The L-amino acid oxidases of snake venom. III. Reversible interactivation of L-amino acid oxidases. Arch. Biochem. Biophys. 33: 377, 1951.
21. Zeller, E. A. Über Phosphatasen. Über eine neue Adenosintri phosphatase. Helvetica Chimica Acta, 33: 821, 1950.
22. Slotta, K. and A. Ballester. Determinacao colorimetrica da hialuronidase dos venenos ofidicos. Mem. Inst. Butantan, 26: 311, 1954.
23. Maeno, H., M. Morimura, S. Mitsuhashi, Y. Sawai and T. Okonogi. Studies on Habu snake venom. 2b. Further purification and enzymic and biological activities of Ha-proteinase. Jap. J. Microbiol. 3: 277, 1959.
24. Maeno, H., S. Mitsuhashi and R. Sato. Studies of Habu snake venom. 2c. Studies on HS-proteinase of Habu venom. Jap. J. Microbiol. 4: 173, 1960.

25. Maeno, H. and S. Mitsuhashi. Studies on Habu snake venom. IV. Fractionation of Habu snake venom by chromatography on CM-cellulose. J. Biochem. 50: 434, 1961.
26. Maeno, H., S. Mitsuhashi, T. Okonogi, S. Hoshi and M. Homma. Studies on Habu snake venom. V. Myolysis caused by phospholipase A in Habu snake venom. Jap. J. Exptl. Med. 32: 55, 1962.
27. Russell, F. E., F. W. Buess and M. Y. Woo. Zootoxico-logical properties of venom phosphodiesterase. Toxicon, 1: 99, 1963.
28. Mandelbaum, F. R. and O. B. Henriques. Purification and properties of Bothrops protease A. Arch. Biochem. Biophys. 104: 369, 1964.
29. Jansen, E. F., M. -D.F. Nutting, R. Jang and A. K. Balls. Inhibition of the proteinase and esterase activities of trypsin and chymotrypsin by diisopropylfluorophosphate: crystallization of inhibited chymotrypsin. J. Biol. Chem. 179: 189, 1949.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

US Army Medical Research Laboratory
Fort Knox, Kentucky 40121

2a. REPORT SECURITY CLASSIFICATION

UNCLASSIFIED

2b. GROUP

3. REPORT TITLE THE EFFECT OF pH, TEMPERATURE, ANTIVENIN AND FUNCTIONAL GROUP INHIBITORS ON THE TOXICITY AND ENZYMATIC ACTIVITIES OF CROTALUS ATROX VENOM

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

5. AUTHOR(S) (Last name, first name, initial)

Brown, John H.

6. REPORT DATE

20 September 1965

7a. TOTAL NO. OF PAGES

13

7b. NO. OF REFS

29

8a. CONTRACT OR GRANT NO.

a. PROJECT NO. 3A014501B71P

c. Task No. 11 (1964)

d. Subtask No. 42

8a. ORIGINATOR'S REPORT NUMBER(S)

640

8b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

9. AVAILABILITY/LIMITATION NOTICES

Qualified requesters may obtain copies of this report from DDC.

10. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

US Army Medical Research and Development Command, Washington, D. C. 20315

3. ABSTRACT

Venom of Crotalus atrox can be detoxified with N-bromosuccinimide. The toxin(s) are not affected by treatment at low pH but are inactivated by heating at 75° or 100°C. At these temperatures all of the enzymes tested are inactivated except phospholipase A.

D FORM 1 JAN 64 1473

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Venoms, Biochemistry						
Enzymes						
Toxins and Antitoxins						
Temperature and Enzyme Activity						
Temperature and Toxicity						
pH and Toxicity						
Antivenin and Enzyme Activity						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).
10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:
 - (1) "Qualified requesters may obtain copies of this report from DDC."
 - (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
 - (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
 - (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
 - (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.
11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.
12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.
14. **KEY WORDS:** Key words are technically meaningful term or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

UNCLASSIFIED

Security Classification